

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶ : C11D 3/22, 3/37, 17/00	A1	(11) International Publication Number: WO 96/37595 (43) International Publication Date: 28 November 1996 (28.11.96)
(21) International Application Number: PCT/US96/07510 (22) International Filing Date: 23 May 1996 (23.05.96) (30) Priority Data: 9510832.0 27 May 1995 (27.05.95) GB 9513404.5 30 June 1995 (30.06.95) GB (71) Applicant (for all designated States except US): THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): FRANKENBACH, Gayle, Marie [US/US]; 10010 Voyager Lane, Cincinnati, OH 45252 (US). PHIPPS, Nicola, Jacqueline [GB/GB]; 19 Wyvern Place, Green Lane, Surrey KT15 2UD (GB). RICHARDSON, Wendy, Victoria, J. [GB/GB]; 70 Edgell Road, Staines, Middlesex TW18 2ES (GB). (74) Agents: REED, T., David et al.; The Procter & Gamble Company; 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).		(81) Designated States: CN, MX, SG, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: AQUEOUS PERSONAL CLEANSING COMPOSITIONS COMPRISING SPECIFIC NONOCCLUSIVE LIQUID POLYOL FATTY ACID POLYESTER (57) Abstract <p>A personal cleansing composition comprising: (a) from about 1 % to about 30 % by weight of a dispersed oil phase comprising at least one nonocclusive liquid polyol fatty acid polyester containing at least four fatty acid ester groups wherein the polyol moiety is selected from sugars and sugar alcohols containing from about 4 to about 8 hydroxyl groups, and wherein each carboxylic acid moiety has from about 8 to about 22 carbon atoms and wherein the liquid polyol fatty acid polyester has a complete melting point of less than about 30 °C; and (b) from about 5 % to about 30 % by weight of water-soluble surfactant selected from anionic, nonionic, zwitterionic and amphoteric surfactants and mixtures thereof. The products provide excellent in-use and efficacy benefits including excellent skin feel during and after use, good lather volume and stability characteristics, skin conditioning, mildness, stability, cleansing, good water-feel and appearance.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

WO 96/37595

PCT/US96/07510

1

AQUEOUS PERSONAL CLEANSING COMPOSITIONS COMPRISING SPECIFIC NONOCCLUSIVE LIQUID POLYOL FATTY ACID POLYESTER.

TECHNICAL FIELD

The present invention relates to cleansing compositions. In particular it relates to mild personal cleansing compositions with good skin feel attributes and foaming properties suitable for simultaneously cleansing and conditioning the skin and/or the hair and which may be used, for example, in the form of foam bath preparations, shower products, skin cleansers, hand, face and body cleansers, shampoos, etc.

BACKGROUND OF THE INVENTION

Mild cosmetic compositions must satisfy a number of criteria including cleansing power, foaming properties and mildness/low irritancy/good feel with respect to the skin, hair and the ocular mucosae. Skin is made up of several layers of cells which coat and protect the keratin and collagen fibrous proteins that form the skeleton of its structure. The outermost of these layers, referred to as the stratum corneum, is known to be composed of 250 Å protein bundles surrounded by 80 Å thick layers. Hair similarly has a protective outer coating enclosing the hair fibre which is called the cuticle. Anionic surfactants can penetrate the stratum corneum membrane and the cuticle and, by delipidization destroy membrane integrity. This interference with skin and hair protective membranes can lead to a rough skin feel and eye irritation and may eventually permit the surfactant to interact with the keratin and hair proteins creating irritation and loss of barrier and water retention functions.

Ideal cosmetic cleansers should cleanse the skin or hair gently, without defatting and/or drying the hair and skin and without irritating the

WO 96/37595

PCT/US96/07510

2

ocular mucosae or leaving skin taut after frequent use. Most lathering soaps, shower and bath products, shampoos and bars fail in this respect.

Certain synthetic surfactants are known to be mild. However, a major drawback of most mild synthetic surfactant systems when formulated for shampooing or personal cleansing is poor lather performance compared to the highest shampoo and bar soap standards. Thus, surfactants that are among the mildest, such as sodium lauryl glyceryl ether sulfonate, (AGS), are marginal in lather. The use of known high sudsing anionic surfactants with lather boosters, on the other hand, can yield acceptable lather volume and quality but at the expense of clinical skin mildness. These two facts make the surfactant selection, the lather and mildness benefit formulation process a delicate balancing act.

Despite the many years of research that have been expended by the toiletries industry on personal cleansing, the broad mass of consumers remain dissatisfied by the mildness of present day cleansing compositions, finding, for example, that they have to apply a separate cosmetic lotion or cream moisturizer to the skin after using a shower or bath preparation in order to maintain skin suppleness and hydration and to counteract the delipidizing effect of the cleanser.

It is known from the art that inclusion of oils in bathing compositions can provide post-use skin feel benefits. However incorporation of oils at levels sufficient to deliver consumer noticeable benefits has until now proved to be a challenge, particularly with respect to the delivery of good, stable lather characteristics in the presence of oil and avoiding unpleasant 'slimy'/'greasy' water feel or oily appearance during use while still delivering a desirable after-use soft skin feel. A further difficulty associated with combining high levels of oil with conventional detergent systems has been the achievement of a stable system in which the oil is deposited onto the skin in a non-greasy manner during use. It has now been found that personal cleansing compositions having excellent skin feel attributes both for in use feel and after use feel as well as good lather volume, stability and feel characteristics in combination with improved product stability can be provided by the use of particular polyol polyester oil components. It has also been found that combination of oil components having particular oil/surfactant

WO 96/37595

PCT/US96/07510

3

solution interfacial tension characteristics are especially valuable for the provision of product stability and skin deposition characteristics

Thus a need exists for personal cleansing products which will not dehydrate the skin or result in loss of skin suppleness, which will provide a level of skin conditioning and emolliency performance which previously has only been provided by a separate post-cleansing cosmetic moisturizer and which will produce a foam which is stable and of high quality both in lather volume and appearance, which are effective hair and skin cleansers, which have good in-use aesthetics, rinsibility characteristics, and which at the same time have stable product and viscosity characteristics and remain fully stable under long term and stressed temperature storage conditions.

SUMMARY OF THE INVENTION

The subject of the present invention is a mild, foam-producing cleansing product suitable for personal cleansing of the skin or hair and which may be used as foam bath and shower products, skin cleansers and shampoos etc. According to one aspect of the invention, there is provided a personal cleansing composition comprising:

- (a) from about 1% to about 30% by weight of a dispersed oil phase comprising a first oil component having an oil/surfactant solution interfacial tension (IFT) of greater than about 1.0 dynes/cm and a second oil component having an oil/surfactant solution interfacial tension (IFT) in the range of from about 0.1 to about 1.0 dynes/cm;
- (b) from about 5% to about 30% by weight of water-soluble surfactant selected from anionic, nonionic, zwitterionic and amphoteric surfactants and mixtures thereof; and
- (c) water;

and wherein the ratio of first to second dispersed oil component is in the range from about 20 : 1 to about 1 : 20.

WO 96/37595

PCT/US96/07510

4

In preferred embodiments, the first oil component is selected from polyol fatty acid polyesters, especially nonocclusive liquid polyol fatty acid polyesters containing at least four fatty acid ester groups and wherein the polyol moiety is selected from sugars and sugar alcohols containing from about 4 to about 8 hydroxyl groups, and wherein each carboxylic acid moiety has from about 8 to about 22 carbon atoms and wherein the liquid polyol fatty acid polyester has a complete melting point of less than about 30°C.

According to a further aspect of the present invention, there is provided an aqueous personal cleansing composition comprising:

- (a) from about 1% to about 30% by weight of a dispersed oil phase comprising at least one nonocclusive liquid polyol fatty acid polyester containing at least four fatty acid ester groups wherein the polyol moiety is selected from sugars and sugar alcohols containing from about 4 to about 8 hydroxyl groups, and wherein each carboxylic acid moiety has from about 8 to about 22 carbon atoms and wherein the liquid polyol fatty acid polyester has a complete melting point of less than about 30°C; and
- (b) from about 5% to about 30% by weight of water-soluble surfactant selected from anionic, nonionic, zwitterionic and amphoteric surfactants and mixtures thereof.

In a highly preferred embodiment, the invention takes the form of a foam producing cleansing composition with excellent lather and stability characteristics, superior skin feel characteristics, improved perceived dryness and assessed tightness and expertly graded dryness, combined with good cleansing ability and conditioning performance.

All concentrations and ratios herein are by weight of the cleansing composition, unless otherwise specified. Surfactant chain lengths are also on a weight average chain length basis, unless otherwise specified.

The cleansing compositions herein are based on a combination of a

WO 96/37595

PCT/US96/07510

5

dispersed oil phase and a water-soluble surfactant system. In preferred compositions the surfactant system comprises an oil dispersing nonionic surfactant and mild auxiliary surfactants, which in general terms can be selected from other nonionic, anionic, amphoteric and zwitterionic surfactants and mixtures thereof. The total level of surfactant, inclusive of anionic, nonionic, zwitterionic, amphoteric and other surfactant components is preferably from about 5% to about 25%, more preferably from about 7% to about 20%, and especially from about 8% to about 16% by weight. The compositions preferably comprise a mixture of

oil dispersing nonionic and anionic surfactants optionally with auxiliary nonionic, zwitterionic and/or amphoteric surfactants. The total level of auxiliary surfactant is in the range from about 0.5% to about 15%, preferably from about 1% to about 10%, more preferably from about 2% to about 6% by weight of the composition, while the level of oil dispersing nonionic surfactant is from about 1% to about 25%, preferably

from about 2% to about 15%, more preferably from about 3% to about 12% by weight and especially from about 4% to about 8%. The weight ratio of anionic surfactant : auxiliary nonionic, zwitterionic and/or amphoteric surfactant is preferably in the range of from about 8:1 to about

1:2. The ratio of auxiliary surfactant to oil dispersing nonionic surfactant is preferably in the range of from 1:10 to about 10:1, more preferably from about 1:5 to about 5:1, most preferably from about 1:2 to about 2:1. The preferred compositions within the scope of the invention comprise mixtures of nonionic surfactants, oil and auxiliary anionic, zwitterionic and/or amphoteric surfactants, wherein the level of dispersed oil phase is from about 3% to about 25%, preferably from about 8% to about 20%, more preferably from about 10% to about 15% by weight of the composition and the weight ratio of oil dispersing nonionic surfactant : dispersed oil phase is in the range of from 1:20 to about 3:2, preferably from about 1:6 to about 1:2, more preferably from about 1:4 to about 1:3.

The dispersed oil phase preferably comprises a mixture of first and second oil components selected on the basis of their oil/surfactant

WO 96/37595

PCT/US96/07510

6

solution interfacial tension characteristics, such combinations being optimum for delivering desirable emulsion stabilisation and skin deposition characteristics.

Oil/surfactant interfacial tension (IFT) measurements indicate the degree to which a surfactant solution can reduce the interfacial tension (IFT) between an oil component and a water phase. IFT measurements are made using a Spinning Drop Interfacial Tensiometer and are taken at 40 - 50°C and 21 Degrees Clark water hardness (428ppm CaCO₃) and 75ppm aqueous surfactant solution concentration. In this test the surfactant system preferred for use is the surfactant system of the final cleansing composition. Where difficulties arise, or for screening purposes a standard aqueous surfactant solution can be used comprising a mixture of ethoxylated alkyl sulphate having two ethoxylated groups and alkyl N-methyl glucose amide in a weight ratio of about 4 : 1. Measurements are undertaken at pH 7 and are reported as the average of the 2, 5 and 10 minute interfacial tension readings.

By 'interfacial tension' (IFT) herein is meant the tension measured at the oil/water interface. IFT measurements using the spinning drop technique, are disclosed by Cafas, Schechter and Wade, "The Measurement of Low Interfacial Tension via the Spinning Drop Technique", ACS Symposium Series No.8 (1975) ADSORPTION AT INTERFACES, beginning at page 234.

The dispersed oil phase suitable for inclusion herein preferably comprises a mixture of a first and a second oil component wherein the first oil component comprises, in preferred embodiments, a polyol polyester and the second oil component comprises one or more oils selected from hydrocarbons, lanolin and lanolin derivatives and animal and vegetable triglycerides. Use of a mixed oil system is valuable both for the effective emulsification of the oils within the product matrix and also for their subsequent deposition upon the skin surface upon product dilution in use. Applicant has found that compositions having mixed first/second oil components as defined deliver improved skin feel both in use and after use versus products containing either the first or second oil components alone.

WO 96/37595

PCT/US96/07510

7

The total level of first and second dispersed oil component present is from about 1% to about 30%, preferably from about 3% to about 25%, more preferably from about 8% to about 20%, most preferably from about 10% to about 15% by weight wherein the weight ratio of first oil component to second oil component is in the range of from about 20 : 1 to about 1 : 20, preferably from about 8 : 1 to about 1 : 8, more preferably from about 4 : 1 to about 1 : 4.

In highly preferred compositions, the dispersed oil phase comprises a nonocclusive liquid or liquifiable polyol fatty acid polyester. Especially preferred are liquid polyol fatty acid polyesters containing at least four fatty acid ester groups and wherein the polyol moiety is selected from sugars and sugar alcohols containing from about 4 to about 8 hydroxyl groups, and wherein each carboxylic acid moiety has from about 8 to about 22 carbon atoms and wherein the liquid polyol fatty acid polyester has a complete melting point of less than about 30°C.

The polyester also preferably has an oil/surfactant interfacial tension (IFT) of greater than about 1.0 dynes/cm, preferably from about 1.1 to about 4.0, more preferably from about 1.2 to about 3.0 and especially from about 1.3 to about 2.0 dynes/cm (as measured under the above standard conditions) and is present at a level of from about 1% to about 10%, preferably from about 1% to about 5%, more preferably from about 2% to about 4% by weight.

The liquid polyol polyesters preferred for use in the present compositions comprise certain polyols, especially sugars or sugar alcohols, esterified with at least four fatty acid groups. Accordingly, preferred polyol starting material should have at least four esterifiable hydroxyl groups. Examples of preferred polyols are sugars, including monosaccharides and disaccharides, and sugar alcohols. Examples of monosaccharides containing four hydroxyl groups are xylose and arabinose and the sugar alcohol derived from xylose, which has five hydroxyl groups, i.e., xylitol. The monosaccharide, erythrose, is not preferred in the practice of this invention since it only contains three hydroxyl groups, but the sugar alcohol derived from erythrose, i.e., erythritol, contains four hydroxyl groups and accordingly can be used. Suitable five hydroxyl group-containing monosaccharides are galactose, fructose, and sorbose. Sugar

PCT/US96/07510

8

Preferred polyols for preparing the polyesters for use in the present invention are selected from the group consisting of erythritol, xylitol, sorbitol, glucose, and sucrose. Sucrose is especially preferred.

The polyol fatty acid polyesters useful in this invention preferably contain at least four fatty acid ester groups. It is not necessary that all of the hydroxyl groups of the polyol be esterified with fatty acid, but it is preferable that the polyester contain no more than two unesterified hydroxyl groups. Most preferably, substantially all of the hydroxyl groups of the polyol are esterified with fatty acid, i.e., the polyol moiety is substantially completely esterified. The fatty acids esterified to the polyol molecule can be the same or mixed, but as noted above, a substantial amount of the unsaturated acid ester groups must be present to provide liquidity.

To illustrate the above points, a sucrose tetra-fatty acid ester would be suitable for use herein, but is not preferred because it has more than two unesterified hydroxyl groups. A sucrose hexa-fatty acid ester would be preferred because it has no more than two unesterified hydroxyl groups. Highly preferred compounds in which all the hydroxyl groups are

WO 96/37595

PCT/US96/07510

9

esterified with fatty acids include the liquid sucrose octa-substituted fatty acid esters.

The following are non-limiting examples of specific polyol fatty acid polyesters containing at least four fatty acid ester groups preferred for use in the present invention: glucose tetraoleate, the glucose tetraesters of soybean oil fatty acids (unsaturated), the mannose tetraesters of mixed soybean oil fatty acids, the galactose tetraesters of oleic acid, the arabinose tetraesters of linoleic acid, xylose tetralinoleate, galactose pentaoleate, sorbitol tetraoleate, the sorbitol hexaesters of unsaturated soybean oil fatty acids, xylitol pentaoleate, sucrose tetraoleate, sucrose pentaoleate, sucrose hexaoleate, sucrose heptaoleate, sucrose octaoleate, and mixtures thereof.

As noted above, highly preferred polyol fatty acid esters are those wherein the fatty acids contain from about 14 to about 18 carbon atoms.

The preferred liquid polyol polyesters preferred for use herein have complete melting points below about 30°C, preferably below about 27.5°C, more preferably below about 25°C. Complete melting points reported herein are measured by Differential Scanning Calorimetry (DSC).

The polyol fatty acid polyesters suitable for use herein can be prepared by a variety of methods well known to those skilled in the art. These methods include: transesterification of the polyol with methyl, ethyl or glycerol fatty acid esters using a variety of catalysts; acylation of the polyol with a fatty acid chloride; acylation of the polyol with a fatty acid anhydride; and acylation of the polyol with a fatty acid, per se. See US-A-2,831,854; US-A-4,005,196, to Jandacek, issued January 25, 1977; US-A-4,005,196, to Jandacek, issued January 25, 1977.

The preferred first oil component for the compositions according to the present invention is a liquid sucrose octa-substituted fatty acid ester.

The compositions of the present invention can also comprise further oil components in the dispersed oil phase. The preferred further oil components are non-polar oils. Highly preferred second oil components have an interfacial tension (IFT) in the range of from about 0.1 to about

WO 96/37595

PCT/US96/07510

10

1.0 dynes/cm, preferably from about 0.2 to about 0.9, more preferably from about 0.3 to about 0.7 dynes/cm (as measured using the above standard conditions) and are present at a level of from about 1% to about 20%, preferably from about 5% to about 15%, more preferably from about 8% to about 15% by weight.

Additional oil components of the dispersed oil phase suitable for use herein include hydrocarbons, lanolin and animal and vegetable triglycerides such as mineral oils, petrolatum and squalene, fatty sorbitan esters (see US-A-3988255, Seiden, issued October 26th 1976), lanolin and oil-like lanolin derivatives, water-insoluble silicones inclusive of non-volatile polyalkyl and polyaryl siloxane gums and fluids, volatile cyclic and linear polyalkylsiloxanes, polyalkoxylated silicones, amino and quaternary ammonium modified silicones, rigid cross-linked and reinforced silicones and mixtures thereof, C₁-C₂₄ esters of C₈-C₃₀ fatty acids such as isopropyl myristate and cetyl ricinoleate, beeswax, saturated and unsaturated fatty alcohols such as behenyl alcohol, almond oil, peanut oil, wheat germ oil, linseed oil, jojoba oil, oil of apricot pits, walnuts, palm nuts, pistachio nuts, sesame seeds, rapeseed, cade oil, corn oil, peach pit oil, poppyseed oil, pine oil, castor oil, soybean oil, avocado oil, safflower oil, coconut oil, hazelnut oil, olive oil, grapeseed oil, and sunflower seed oil, and C₁-C₂₄ esters of dimer and trimer acids such as diisopropyl dimerate, diisostearylmalate, diisostearyldimerate and triisostearyltrimerate and mixtures thereof.

The most preferred second oil components are non-polar oils selected from mineral oil, petrolatum, water-insoluble silicones, soya bean oil and the like and mixtures thereof. Especially preferred for use herein is mineral oil.

Preferred compositions include an oil dispersing nonionic surfactant at a level of from about 1% to about 25%, preferably from about 2% to about 15%, more preferably from about 3% to about 12%, most preferably from about 4% to about 8% by weight.

Oil dispersing nonionic surfactants suitable for inclusion in the compositions according to the present invention have excellent oil

WO 96/37595

PCT/US96/07510

11


dispersing characteristics as demonstrated by microscopy and Immersion tests.

To demonstrate the suitability of a nonionic surfactant for incorporation in the compositions according to the present invention a test matrix including the test nonionic and all other composition ingredients is prepared. A portion of this prototype is then viewed under a microscope using a Nikon Optipot-2 Videomicrowatcher. Surfactants suitable for incorporation in the compositions according to the present invention typically produce homogeneous (opaque) product matrices in which the oil droplets have a mean diameter in the range of from about 1 microns to about 30 microns, preferably from about 2 microns to about 20 microns.

Highly preferred dispersed oil systems suitable for the compositions according to the present invention have an average oil droplet size of from about 3 microns to about 10 microns as measured by a Nikon Optipot-2 Videomicrowatcher.

In the Immersion Test, the in-use characteristics of prototype matrices are assessed. In essence, 6 ml of prototype matrix is dispersed in 20 litres of water at 40°C. The physical appearance of the resulting solution is then assessed. Compositions which result in turbid (cloudy) solutions wherein no oil droplets are visible to the naked eye are regarded as demonstrating the desired behaviour, providing they additionally deliver the required skin feel.

Oil dispersing nonionic surfactants suitable for inclusion in the compositions according to the present invention are selected from C₁₂-C₁₈ polyhydroxy fatty acid amide surfactants, preferably C₁₂-C₁₆ polyhydroxy fatty acid amide surfactants, more preferably C₁₂-C₁₄ polyhydroxy fatty acid amide surfactants having the general formula (I).



The preferred N-alkyl, N-alkoxy or N-aryloxy, polyhydroxy fatty acid amide surfactants according to formula (I) are those in which R⁸ is C₅-

WO 96/37595

PCT/US96/07510

12

C₃₁ hydrocarbonyl, preferably C₉-C₁₇ hydrocarbonyl, including straight-chain and branched chain alkyl and alkenyl, or mixtures thereof and R₉ is typically C₁-C₈ alkyl or hydroxyalkyl, preferably methyl, or a group of formula -R¹-O-R² wherein R¹ is C₂-C₈ hydrocarbonyl including straight-chain, branched-chain and cyclic (including aryl), and is preferably C₂-C₄ alkylene, R² is C₁-C₈ straight-chain, branched-chain and cyclic hydrocarbonyl including aryl and oxyhydrocarbonyl, and is preferably C₁-C₄ alkyl, especially methyl, or phenyl. Z₂ is a polyhydroxyhydrocarbonyl moiety having a linear hydrocarbonyl chain with at least 2 (in the case of glyceraldehyde) or at least 3 hydroxyls (in the case of other reducing sugars) directly connected to the chain, or an alkoxylated derivative (preferably ethoxylated or propoxylated) thereof. Z₂ preferably will be derived from a reducing sugar in a reductive amination reaction, most preferably Z₂ is a glycidyl moiety. Suitable reducing sugars include glucose, fructose, maltose, lactose, galactose, mannose, and xylose, as well as glyceraldehyde. As raw materials, high dextrose corn syrup, high fructose corn syrup, and high maltose corn syrup can be utilized as well as the individual sugars listed above. These corn syrups may yield a mix of sugar components for Z₂. It should be understood that it is by no means intended to exclude other suitable raw materials. Z₂ preferably will be selected from the group consisting of -CH₂-(CHOH)_n-CH₂OH, -CH(CH₂OH)-(CHOH)_{n-1}-CH₂OH, CH₂(CHOH)₂(CHOR')CHOH-CH₂OH, where n is an integer from 1 to 5, inclusive, and R' is H or a cyclic mono- or poly-saccharide, and alkoxylated derivatives thereof. As noted, most preferred are glycidyls wherein n is 4, particularly -CH₂-(CHOH)₄-CH₂OH.

In compounds of the above formula, R_g-CO-N< can be, for example, cocoamide, stearamide, oleamide, lauramide, myristamide, capricamide, palmitamide, tallowamide, etc.

A preferred process for making the above compounds having formula (I) comprises reacting a fatty acid triglyceride with an N-substituted polyhydroxy amine in the substantial absence of lower (C₁-C₄) alcoholic solvent, but preferably with an alkoxylated alcohol or alkoxylated alkyl phenol such as NEODOL and using an alkoxide catalyst at temperatures of from about 50°C to about 140°C to provide high yields (90-98%) of the desired products. Suitable processes for making the desired polyhydroxy

fatty acid amide materials are outlined in US-A-5,194,639 and US-A-5,380,891.

The most preferred polyhydroxy fatty acid amide has the formula $R_8(CO)N(CH_3)CH_2(CHOH)_4CH_2OH$ wherein R_8 is a C11-C17 straight chain alkyl or alkenyl group.

In preferred compositions the surfactant system comprises an oil dispersing nonionic surfactant and auxiliary surfactants wherein the auxiliary surfactant is selected from anionic, zwitterionic, amphoteric and auxiliary nonionic surfactants and mixtures thereof. The level of auxiliary surfactant is from about 0.5% to about 15%, preferably from about 1% to about 10%, more preferably from about 2% to about 6% by weight.

Anionic surfactants suitable for inclusion in the compositions of the invention can generally be described as mild synthetic detergent surfactants and include ethoxylated alkyl sulfates, alkyl glyceryl ether sulfonates, methyl acyl taurates, fatty acyl glycinate, N-acyl glutamates, acyl isethionates, alkyl sulfosuccinates, alpha-sulfonated fatty acids, their salts and/or their esters, alkyl ethoxy carboxylates, alkyl phosphate esters, ethoxylated alkyl phosphate esters, acyl sarcosinates and fatty acid/protein condensates, and mixtures thereof. Alkyl and/or acyl chain lengths for these surfactants are C₈-C₂₂, preferably C₁₀-C₁₈.

Preferred for use herein from the viewpoint of optimum mildness and lathering characteristics are the salts of sulfuric acid esters of the reaction product of 1 mole of a higher fatty alcohol and from about 1 to about 12 moles of ethylene oxide, with sodium and magnesium being the preferred counterions. Particularly preferred are the alkyl sulfates containing from about 2 to 6, preferably 2 to 4 moles of ethylene oxide, such as sodium laureth-2 sulfate, sodium laureth-3 sulfate and magnesium sodium laureth-3.6 sulfate. In preferred embodiments, the anionic surfactant contains at least about 50%, especially at least about 75% by weight of ethoxylated alkyl sulfate.


WO 96/37595

PCT/US96/07510


14

The compositions for use herein suitably also contain an amphoteric surfactant. Amphoteric surfactants suitable for use in the compositions of the invention include:

- (a) imidazolinium surfactants of formula (II)



wherein R₁ is C₇-C₂₂ alkyl or alkenyl, R₂ is hydrogen or CH₂Z, each Z is independently CO₂M or CH₂CO₂M, and M is H, alkali metal, alkaline earth metal, ammonium or alkanolammonium; and/or ammonium derivatives of formula (III)



wherein R₁, R₂ and Z are as defined above;

- (b) aminoalkanoates of formula (IV)



and iminodialkanoates of formula (V)



wherein n and m are numbers from 1 to 4, and R₁ and M are independently selected from the groups specified above; and

- (c) mixtures thereof.

WO 96/37595

PCT/US96/07510

15

Suitable amphoteric surfactants of type (a) are marketed under the trade name Miranol and Empigen and are understood to comprise a complex mixture of species. Traditionally, the Miranols have been described as having the general formula II, although the CTFA Cosmetic Ingredient Dictionary, 4th Edition indicates the non-cyclic structure III. In practice, a complex mixture of cyclic and non-cyclic species is likely to exist and both definitions are given here for sake of completeness. Preferred for use herein, however, are the non-cyclic species.

Examples of suitable amphoteric surfactants of type (a) include compounds of formula II and/or III in which R_1 is C_8H_{17} (especially iso-capryl), C_9H_{19} and $C_{11}H_{23}$ alkyl. Especially preferred are the compounds in which R_1 is C_9H_{19} , Z is CO_2M and R_2 is H; the compounds in which R_1 is $C_{11}H_{23}$, Z is CO_2M and R_2 is CH_2CO_2M ; and the compounds in which R_1 is $C_{11}H_{23}$, Z is CO_2M and R_2 is H.

In CTFA nomenclature, materials preferred for use in the present invention include cocoamphocarboxypropionate, cocoamphocarboxypropionic acid, and especially cocoamphoacetate and cocoamphodiacetate (otherwise referred to as cocoamphocarboxyglycinate). Specific commercial products include those sold under the trade names of Empigen CDL60 and CDR 60 (Albright & Wilson), Miranol H2M Conc. Miranol C2M Conc. N.P., Miranol C2M Conc. O.P., Miranol C2M SF, Miranol CM Special (Rhône-Poulenc); Alkateric 2CIB (Alkaril Chemicals); Amphoterger W-2 (Lonza, Inc.); Monateric CDX-38, Monateric CSH-32 (Mona Industries); Rewoteric AM-2C (Rewo Chemical Group); and Schercotic MS-2 (Scher Chemicals).

It will be understood that a number of commercially-available amphoteric surfactants of this type are manufactured and sold in the form of electroneutral complexes with, for example, hydroxide counterions or with anionic sulfate or sulfonate surfactants, especially those of the sulfated C_8 - C_{18} alcohol, C_8 - C_{18} ethoxylated alcohol or C_8 - C_{18} acyl glyceride types. Preferred from the viewpoint of mildness and product stability, however, are compositions which are essentially free of (non-ethoxylated) sulfated alcohol surfactants. Note also that the

WO 96/37595


PCT/US96/07510

16

concentrations and weight ratios of the amphoteric surfactants are based herein on the uncomplexed forms of the surfactants, any anionic surfactant counterions being considered as part of the overall anionic surfactant component content.

Examples of suitable amphoteric surfactants of type (b) include salts, especially the triethanolammonium salts and salts of N-lauryl-beta-amino propionic acid and N-lauryl-imino-dipropionic acid. Such materials are sold under the trade name Deriphat by Henkel and Mirataine by Rhône-Poulenc. Amphoterics preferred for use herein, however, are those of formula II and/or III.

The compositions of the invention may also include auxiliary nonionic surfactants. Suitable auxiliary nonionic surfactants for use herein can be selected from C₁₂-C₁₄ fatty acid mono- and diethanolamides and sucrose polyester surfactants, water soluble vegetable and animal-derived emollients (oil derived) such as triglycerides with a polyglycol chain inserted; ethoxylated mono and di-glycerides, polyethoxylated lanolins and shea butter derivatives and mixtures thereof. One preferred class of oil-derived auxiliary nonionic surfactants for use herein have the general formula (VI)



wherein n is from about 5 to about 200, preferably from about 20 to about 100, more preferably from about 30 to about 85, and wherein R comprises an aliphatic radical having on average from about 5 to 20 carbon atoms, preferably from about 9 to 18 carbon atoms.

Suitable ethoxylated oils and fats of this class include polyethyleneglycol derivatives of glyceryl cocoate, glyceryl caproate, glyceryl caprylate, glyceryl tallowate, glyceryl palmate, glyceryl stearate, glyceryl laurate, glyceryl oleate, glyceryl ricinoleate, and glyceryl fatty esters derived from triglycerides, such as palm oil, almond oil, and corn oil, preferably glyceryl tallowate and glyceryl cocoate.

WO 96/37595

PCT/US96/07510

17

Suitable oil derived auxiliary nonionic surfactants of this class are available from Croda Inc. (New York, USA) under their Crovol line of materials such as Crovol EP40 (PEG 20 evening primrose glyceride), Crovol EP 70 (PEG 60 evening primrose glyceride) Crovol A-40 (PEG 20 almond glyceride), Crovol A-70 (PEG 60 almond glyceride), Crovol M-40 (PEG 20 maize glyceride), Crovol M-70 (PEG 60 maize glyceride), Crovol PK-40 (PEG 12 palm kernel glyceride), and Crovol PK-70 (PEG 45 palm kernel glyceride) and under their Solan range of materials such as Solan E, E50 and X polyethoxylated lanolins. Also especially preferred herein is PEG (6) capric/caprylic glyceride (Softigen 767). Further suitable surfactants of this class are commercially available from Sherex Chemical Co. (Dublin, Ohio, USA) under their Varonic LI line of surfactants. These include, for example, Varonic LI 48 (polyethylene glycol (n=80) glyceryl tallowate, alternatively referred to as PEG 80 glyceryl tallowate), Varonic LI 2 (PEG 28 glyceryl tallowate), Varonic LI 420 (PEG 200 glyceryl tallowate), and Varonic LI 63 and 67 (PEG 30 and PEG 80 glyceryl cocoates). Other water soluble vegetable-derived emollients suitable for use are PEG derivatives of corn, avocado and babassu oil.

Also suitable for use herein are auxiliary nonionic surfactants derived from composite vegetable fats extracted from the fruit of the Shea Tree (*Butyrospermum Karkii* Kotschy) and derivatives thereof. This vegetable fat, known as Shea Butter is widely used in Central Africa for a variety of means such as soap making and as a barrier cream, it is marketed by Sederma (78610 Le Perray En Yvelines, France). Also of interest are ethoxylated derivatives of Shea butter available from Karlshamn Chemical Co. (Columbos, Ohio, USA) under their Lipex range of chemicals, such as Lipex 102 E-75 (ethoxylated mono, di-glycerides of Shea butter) and from Croda Inc. (New York) under their Crovol line of materials, such as Crovol SB70 (RTM) (ethoxylated Shea butter). Similarly, ethoxylated derivatives of Mango, Cocoa and Illipe butter may be used in compositions according to the invention. Although these are classified as ethoxylated nonionic surfactants it is understood that a certain proportion may remain as non-ethoxylated vegetable oil or fat.

WO 96/37595

18


PCT/US96/07510

Other suitable auxiliary oil-derived nonionic surfactants include ethoxylated derivatives of almond oil, peanut oil, wheat germ oil, linseed oil, jojoba oil, oil of apricot pits, walnuts, palm nuts, pistachio nuts, sesame seeds, rapeseed, cade oil, corn oil, peach pit oil, poppyseed oil, pine oil, castor oil, soybean oil, avocado oil, safflower oil, coconut oil, hazelnut oil, olive oil, grapeseed oil, and sunflower seed oil.

Oil derived auxiliary nonionic surfactants highly preferred for use herein from the viewpoint of optimum mildness and skin feel characteristics are PEG 60 evening primrose triglycerides; PEG 55 lanolin polyethoxylated derivatives and ethoxylated derivatives of Shea butter.


The compositions herein preferably also contain a zwitterionic surfactant.

Betaine surfactants suitable for inclusion in the composition of the invention include alkyl betaines of the formula $R_5R_6R_7N^+(CH_2)_nM$ (VII) and amido betaines of the formula (VIII)



wherein R_5 is C_{11} - C_{22} alkyl or alkenyl, R_6 and R_7 are independently C_1 - C_3 alkyl, M is H, alkali metal, alkaline earth metal, ammonium or alkanolammonium, and n , m are each numbers from 1 to 4. Preferred betaines include cocoamidopropyldimethylcarboxymethyl betaine, laurylamidopropyldimethylcarboxymethyl betaine and Tego betaine (RTM).

Water-soluble auxiliary sultaine surfactants suitable for inclusion in the compositions of the present invention include alkyl sultaines of the formula (IX);




WO 96/37595

PCT/US96/07510

19

wherein R₁ is C₇ to C₂₂ alkyl or alkenyl, R₂ and R₃ are independantly C₁ to C₃ alkyl, M is H, alkali metal, alkaline earth metal, ammonium or alkanolammonium and m and n are numbers from 1 to 4. Preferred for use herein is coco amido propylhydroxy sultaine.

Water-soluble auxiliary amine oxide surfactants suitable for inclusion in the compositions of the present invention include alkyl amine oxide R₅R₆R₇NO and amido amine oxides of the formula (X)



wherein R₅ is C₁₁ to C₂₂ alkyl or alkenyl, R₆ and R₇ are independantly C₁ to C₃ alkyl, M is H, alkali metal, alkaline earth metal, ammonium or alkanolammonium and m is a number from 1 to 4. Preferred amine oxides include cocoamidopropylamine oxide, lauryl dimethyl amine oxide and myristyl dimethyl amine oxide.

The compositions of the invention may also contain a cationic or nonionic polymeric skin or hair conditioning agent at a level from about 0.01 % to about 5 %, preferably from about 0.04 % to about 2 % and especially from about 0.05 % to about 1 % by weight. The polymer is found to be valuable for enhancing the creaminess and quality of the foam as well as providing a hair or skin conditioning utility.

Suitable polymers are high molecular weight materials (mass-average molecular weight determined, for instance, by light scattering, being generally from about 2,000 to about 3,000,000, preferably from about 5,000 to about 1,000,000).

Useful polymers are the cationic, nonionic, amphoteric, and anionic polymers useful in the cosmetic field. Preferred are cationic and

WO 96/37595

PCT/US96/07510

20

nonionic polymers used in the cosmetic fields as hair or skin conditioning agents.

Representative classes of polymers include cationic and nonionic polysaccharides; cationic and nonionic homopolymers and copolymers derived from acrylic and/or methacrylic acid; cationic and nonionic cellulose resins; cationic copolymers of dimethyldiallylammonium chloride and acrylic acid; cationic homopolymers of dimethyldiallylammonium chloride; cationic polyalkylene and ethoxypolyalkylene imines; quaternized silicones, and mixtures thereof.

By way of exemplification, cationic polymers suitable for use herein include cationic guar gums such as hydroxypropyl trimethyl ammonium guar gum (d.s. of from 0.11 to 0.22) available commercially under the trade names Jaguar C-14-S(RTM) and Jaguar C-17(RTM) and also Jaguar C-16(RTM), which contains hydroxypropyl substituents (d.s. of from 0.8-1.1) in addition to the above-specified cationic groups, and quaternized cellulose ethers available commercially under the trade names Ucare Polymer JR and Celquat. Other suitable cationic polymers are homopolymers of dimethyldiallylammonium chloride available commercially under the trade name Merquat 100, copolymers of dimethyl aminoethylmethacrylate and acrylamide, copolymers of dimethyldiallylammonium chloride and acrylamide, available commercially under the trade names Merquat 550 and Merquat S, quaternized vinyl pyrrolidone acrylate or methacrylate copolymers of amino alcohol available commercially under the trade name Gafquat, and polyalkyleneamines such as polyethylenimine and ethoxylated polyethylenimine.

Nonionic polymers suitable for use in the compositions according to the present invention include any conventionally used nonionic polymer and preferably those of the Pluronic and Synperonic group of polyoxyethylene polyoxypropylene block copolymers available from BASF, such as Pluronic L-121 and ICI, such as synperonic PE-F127.

Anionic polymers suitable herein include hydrophobically-modified cross-linked polymers of acrylic acid having amphipathic properties as marketed by B F Goodrich under the trade name Pemulen TRI and

WO 96/37595

PCT/US96/07510

21

Pemulen TR2; and the carboxyvinyl polymers sold by B F Goodrich under the trade mark Carbopol and which consist of polymers of acrylic acid cross-linked with polyallyl sucrose or polyallyl pentaerythritol, for example, Carbopol 934, 940 and 950.

The viscosity of the final composition (Helipath, Spindle A, 10 rpm, 25°C, neat) is preferably at least about 500 cps, more preferably from about 1,000 to about 10,000 cps, especially from about 1,000 to about 5,000 cps.

The cleansing compositions can optionally include a hair or skin moisturizer which is soluble in the cleansing composition matrix. The preferred level of moisturizer is from about 0.5% to about 20% by weight. In preferred embodiments, the moisturizer is selected from:


1. water-soluble liquid polyols;
2. essential amino acid compounds found naturally occurring in the stratum corneum of the skin; and
3. water-soluble nonpolyol nonocclusives and mixtures thereof.

Some examples of more preferred nonocclusive moisturizers are glycerine, polyethylene glycol, propylene glycol, sorbitol, polyethylene glycol and propylene glycol ethers of methyl glucose (e.g. methyl gluceth-20), polyethylene glycol and propylene glycol ethers of lanolin alcohol (e.g. Solulan-75), sodium pyrrolidone carboxylic acid, lactic acid, urea, L-proline, guanidine, pyrrolidone, hydrolyzed protein and other collagen-derived proteins, aloe vera gel and acetamide MEA and mixtures thereof. Of the above, glycerine is highly preferred.

An additional optional component of the composition of the invention is an adduct prepared from vegetable oils containing non-conjugated polyunsaturated fatty acid esters which are conjugated and oxidized and then modified via Diels-Alder addition with a member of the group consisting of acrylic acid, fumaric acid and maleic anhydride. The vegetable oil adduct preferably has the general formula (X).

WO 96/37595

PCT/US96/07510



wherein x , y are integers of from 3 to 9, R_3 and R_4 are independently selected from saturated and unsaturated C_7 - C_{22} hydrocarbonyl, each Z being CO_2M and wherein M is H , or a salt forming cation, preferably alkalimetal, ammonium or alkanol ammonium. The adducts and their preparation are described in US-A-4740367, the adducts being marketed under the trade name Ceraphyl GA (Van Dyke). The vegetable oil adduct is preferably added from about 0.01 % to about 5 %, preferably from about 0.05 % to about 2 %, more preferably from about 0.1 % to about 1 % by weight of the composition.

The compositions according to the present invention can also include a stabilising system. Preferred stabilising systems suitable for inclusion in the compositions according to the present invention comprise a primary and secondary stabilising agent wherein the primary stabilising agent is an optionally modified clay or clay like material and the secondary stabilising agent is a hetero polysaccharide gum. Use of a mixed stabilising system is valuable for the delivery of lather having good volume, stability and appearance characteristics in combination with improvements in skin feel.

The total level of primary and secondary stabilising agent present is from about 0.01 % to about 15 %, preferably from about 0.05 % to about 12 %, more preferably from about 0.1 % to about 10 % most preferably from about 0.5 % to about 5 % by weight wherein the ratio of primary stabiliser to secondary stabiliser is in the range of from about 32:1 to about 1:1, preferably from about 16:1 to about 1:1, more preferably from about 4:1 to about 2:1.

In preferred compositions according to the invention the primary stabiliser is a modified clay based material.

Suitable primary suspending agents for the compositions of the present invention include magnesium aluminium silicate ($Al_2Mg_8Si_2$), bentonite, hectorite and derivatives thereof. Magnesium aluminium

The preferred secondary stabilising agent in the compositions according to the invention is xanthan gum (xanthan / corn sugar gum) which is a heteropolysaccharide gum produced by a pure-culture fermentation of a carbohydrate with *Xanthomonas campestris* having a molecular weight of greater than about 1,000,000. It is believed to contain D-glucose, D-mannose and D-glucuronate in the molar ratios of 2.8 : 2.0 : 2.0. The polysaccharide is partially acetylated with 4.7% acetyl. This biosynthetic gum material is commercially available from Calgon under the trade name Kelgum CG (RTM) and from Kelco (a division of Merck & Co., Inc.) , Meer and Vanderbilt under the respective trade names Keltrol (RTM), Merezan 8 (RTM) and Rhodigel (RTM) as well as from a variety of other sources. Xanthan gum mixtures are also available from Calgon, Alban Muller and others and are also suitable for inclusion in the compositions of the present invention. Further information on xanthan gum is to be found in Whistler, Roy L. (Editor) Industrial Gums - Polysaccharides and Their Derivatives New York: Academic Press, 1973.

WO 96/37595

PCT/US96/07510

24

The compositions according to the present invention can also contain hydrotropes to impart improved low temperature stability characteristics. A suitable low temperature hydrotrope for the present compositions is sodium C₈ alkyl sulphate.

A number of additional optional materials can be added to the cleansing compositions. Such materials include proteins and polypeptides and derivatives thereof; water-solubilizable preservatives such as DMDM Hydantoin, Germall 115, methyl, ethyl, propyl and butyl esters of hydroxybenzoic acid, EDTA, Euxyl (RTM) K400, Bronopol (2-bromo-2-nitropropane-1,3-diol), sodium benzoate, potassium sorbate and 2-phenoxyethanol; other moisturizing agents such as hyaluronic acid, chitin and starch-grafted sodium polyacrylates such as Sanwet (RTM) IM-1000, IM-1500 and IM-2500 available from Celanese Superabsorbent Materials, Portsmouth, VA, USA and described in US-A-4,076,663; solvents such as hexylene glycol and propylene glycol; anti-bacterial agents such as Oxeco (phenoxy isopropanol); low temperature phase modifiers such as ammonium ion sources (e.g. NH₄ Cl); viscosity control agents such as magnesium sulfate, citrate salts and other electrolytes; colouring agents; pearlescers and opacifiers such as styrene PVP polymers such as Lytron 631 (RTM), TiO₂ and TiO₂-coated mica; perfumes and perfume solubilizers; and zeolites such as Valfour BV400 and derivatives thereof and Ca²⁺/Mg²⁺ sequestrants such as polycarboxylates, amino polycarboxylates, polyphosphates, polyphosphonates, amino polyphosphonates and gluconates etc and pH adjusting agents such as citric acid and salts thereof. Water is also present at a level preferably of from about 40% to about 94% preferably at least about 50% by weight of the compositions herein.

The pH of the compositions is preferably from about 4 to about 8.

WO 96/37595

PCT/US96/07510

25

The invention is illustrated by the following non-limiting examples.

In the examples, all concentrations are on a 100% active basis and the abbreviations have the following designation:

Oil 1	Liquid sucrose octaoleate having a complete melting point of less than about 30°C and an IFT of about 1.45 dynes/cm (on a 4:1 anionic : GA aqueous solution basis) wherein IFT is measured as described herein.
Oil 2	Mineral Oil having an IFT of about 0.38 (measured as detailed for Oil 1).
GA	Polyhydroxy fatty acid amide of formula I in which R ₈ is C ₁₁ -C ₁₇ alkyl, R ₉ is methyl, and Z ₂ is CH ₂ (CHOH) ₄ CH ₂ OH
Anionic	Sodium laureth-2 sulfate
Amphoterics	Empigen CDL 60 - an aqueous mixture of 23.5% cocoamphoacetate (in which R ₁ is coconut alkyl R ₂ is H, and Z is CO ₂ Na) and 1.35% cocoamphodiaceate (in which R ₁ is coconut alkyl, R ₂ is CH ₂ CO ₂ Na and Z is CO ₂ Na).
Betaine	Cocoamidopropyldimethylcarboxymethylbetaine
Solan	Solan (RTM) E (PEG 55 lanolin)
Shea	Hydrosoluble Shea Butter (PEG 75)
Preservative	DMDM Hydantoin
Stabiliser	MgAl silicate CMC (Veegum Plus RTM)/Xanthan Gum
Hydrotrope	Sodium C ₈ Alkyl Sulphate

Examples I to VII

	I	II	III	IV	V	VI	VII
GA	2.0	5.0	6.0	4.0	1.5	6.0	5.0
Anionic	4.0	2.0	1.0	2.5	6.0	0.5	1.5
Amphoteric	-	2.0	-	-	-	-	0.5
Betaine	0.5	0.5	1.0	-	-	0.5	0.5
Oil 1	4.0	1.0	5.0	4.0	3.0	1.0	2.0
Oil 2	8.0	9.0	8.0	12.0	10.0	12.0	8.0
Solan	0.5	0.5	-	-	-	-	-
Shea	-	-	-	0.5	-	-	-
Stabiliser	0.8	1.6	4.0	0.3	1.6	2.0	0.2
Hydrotrope	0.1	-	-	1.0	2.0	0.5	0.2
Perfume	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Preservative	0.15	0.15	0.15	0.15	0.15	0.2	0.2
Water	to 100						

WO 96/37595

PCT/US96/07510

27

Compositions I to VII are prepared by first hydrating any Veegum type material as a pre-mix. Next all of the oil materials are separately blended together using by stirring with water at ambient temperature. If xanthan gum type material is present it may be slurried by stirring with this mixture. Separately, the surfactant materials are hydrated along with any additional skin feel agents, preservatives and hydrotropes by mixing with water and heating to between about 20°C and about 90°C. Finally, the oil blend is added to the hydrated Veegum pre-mix (if present) and then the stirred surfactant mixture is added to the combined mixture and this final blend is stirred and cooled to ambient temperature and the remaining water, preservatives, perfume and any other materials are added.

Finally, the compositions have a viscosity (Helipath, Spindle A, 10 rpm, 25°C, neat) in the range from 1,000 to 10,000 cps.

The products provide excellent in-use and efficacy benefits including excellent skin feel during and after use, good lather volume and stability characteristics, skin conditioning, mildness, stability, cleansing, good water-feel and appearance.

WO 96/37595

PCT/US96/07510

28

CLAIMS

1. An aqueous personal cleansing composition comprising:
 - (a) from about 1% to about 30% by weight of a dispersed oil phase comprising at least one nonocclusive liquid polyol fatty acid polyester containing at least four fatty acid ester groups wherein the polyol moiety is selected from sugars and sugar alcohols containing from about 4 to about 8 hydroxyl groups, and wherein each carboxylic acid moiety has from about 8 to about 22 carbon atoms and wherein the liquid polyol fatty acid polyester has a complete melting point of less than about 30°C; and
 - (b) from about 5% to about 30% by weight of water-soluble surfactant selected from anionic, nonionic, zwitterionic and amphoteric surfactants and mixtures thereof.
2. A composition according to Claim 1 wherein the nonocclusive liquid polyol fatty acid polyester has an oil/surfactant solution interfacial tension (IFT) of greater than about 1.0 dynes/cm.
3. A composition according to Claim 1 or 2 wherein the water-soluble surfactant comprises:
 - (a) from about 1% to about 25% by weight of an oil dispersing nonionic surfactant; and
 - (b) from about 0.5% to about 15% by weight of auxiliary surfactant selected from other nonionic, anionic, zwitterionic and/or amphoteric surfactants and mixtures thereof.
4. A composition according to any of Claims 1 to 3 wherein the composition has a viscosity (Helipath, Spindle A, 10 rpm, 25°C, neat) in the range from 1,000 to 10,000 cps .
5. A composition according to any of Claims 1 to 4 wherein the liquid polyol fatty acid polyester is selected from glucose tetraoleate, the

WO 96/37595

PCT/US96/07510

29

glucose tetraesters of soybean oil fatty acids (unsaturated), the mannose tetraesters of mixed soybean oil fatty acids, the galactose tetraesters of oleic acid, the arabinose tetraesters of linoleic acid, xylose tetralinoleate, galactose pentaoleate, sorbitol tetraoleate, the sorbitol hexaesters of unsaturated soybean oil fatty acids, xylitol pentaoleate, sucrose tetraoleate, sucrose pentaoleate, sucrose hexaoleate, sucrose hepatoleate, sucrose octaoleate, and mixtures thereof.


6. A composition according to any of Claims 1 to 5 wherein the liquid polyol fatty acid polyester is sucrose octaoleate.
7. A composition according to any of Claims 1 to 6 wherein the liquid polyol fatty acid polyester is present at a level of from about 1% to about 10%, preferably from about 1% to about 5%, more preferably from about 2% to about 4% by weight.
8. A composition according to any of Claims 1 to 7 wherein the dispersed oil phase additionally comprises one or more oils selected from hydrocarbons, animal and vegetable triglycerides lanolin and oil-like lanolin derivatives, water-insoluble silicones inclusive of non-volatile polyalkyl and polyaryl siloxane gums and fluids, volatile cyclic and linear polyalkylsiloxanes, polyalkoxylated silicones, amino and quaternary ammonium modified silicones, rigid cross-linked and reinforced silicones and mixtures thereof, C₁-C₂₄ esters of C₈-C₃₀ fatty acids such as isopropyl myristate and cetyl ricinoleate, beeswax, saturated and unsaturated fatty alcohols such as behenyl alcohol, almond oil, peanut oil, wheat germ oil, linseed oil, jojoba oil, oil of apricot pits, walnuts, palm nuts, pistachio nuts, sesame seeds, rapeseed, cade oil, corn oil, peach pit oil, poppyseed oil, pine oil, castor oil,
soybean oil, avocado oil, safflower oil, coconut oil, hazelnut oil, olive oil, grapeseed oil, and sunflower seed oil, and C₁-C₂₄ esters
of dimer and trimer acids such as diisopropyl dimerate, diisostearylmalate, diisostearyldimerate and triisostearyltrimerate and mixtures thereof.

WO 96/37595

PCT/US96/07510

30

9. A composition according to Claim 8 wherein the additional oil component comprises one or more oils selected from mineral oil, petrolatum, water insoluble silicones, soya bean oil and mixtures thereof.
10. A composition according to Claims 8 or 9 wherein the additional oil component is present at a level of from about 1% to about 20%, preferably from about 5% to about 15%, most preferably from about 8% to about 15% by weight.
11. A composition according to any of Claims 1 to 10 wherein the total level of oil components is from about 3% to about 25%, preferably from about 8% to about 20%, more preferably from about 10% to about 15%.
12. A composition according to any of Claims 8 to 10 wherein the ratio of liquid polyol fatty acid polyester oil to additional oil component is from about 8:1 to about 1:8, preferably from about 4:1 to about 1:4.
13. A composition according to Claims 3 wherein the level of oil dispersing nonionic surfactant is from about 2% to about 15%, preferably from about 3% to about 12%, more preferably from about 4% to about 8% by weight.
14. A composition according to Claims 3 or 13 wherein the oil dispersing nonionic surfactant is selected from C₁₂-C₁₈ polyhydroxy fatty acid amide surfactants having the general formula (I):



wherein R₉ is C₁₋₄ alkyl, and R₈ is C₇-C₁₉ alkyl or alkenyl, or mixture thereof; and Z₂ is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxyated derivative thereof.

WO 96/37595

PCT/US96/07510

31

15. A composition according to Claim 14 wherein the oil dispersing nonionic surfactant comprises a polyhydroxy fatty acid amide having the formula $R_8(CO)N(CH_3)CH_2(CHOH)_4CH_2OH$ wherein R_8 is a C_{11} - C_{17} straight chain alkyl or alkenyl group.
16. A composition according to Claim 15 wherein the oil dispersing nonionic surfactant comprises a C_{12} - C_{14} polyhydroxy fatty acid amide.
17. A composition according to Claim 3 comprising from about 0.5% to about 10%, more preferably from about 0.5% to about 5%, most preferably from about 1% to about 3.5% by weight in total of auxiliary surfactant.
18. A composition according to Claim 3 wherein the weight ratio of oil dispersing nonionic surfactant to dispersed oil phase is in the range of from 1:20 to about 3:2, preferably from about 1:6 to about 1:2, more preferably from about 1:4 to about 1:3.
19. A composition according to Claim 3 wherein the ratio of auxiliary surfactant to oil dispersing nonionic surfactant is in the range of from 1:10 to about 10:1, preferably from about 1:5 to about 5:1, more preferably from about 1:2 to about 2:1.
20. A composition according to any of Claims 1 to 19 wherein the anionic surfactant is selected from ethoxylated alkyl sulfates, alkyl glyceryl ether sulfonates, methyl acyl taurates, fatty acyl glycinate, alkyl ethoxy carboxylates, N-acyl glutamates, acyl isethionates, alkyl sulfosuccinates, alpha-sulfonated fatty acids, their salts and/or their esters, alkyl phosphate esters, ethoxylated alkyl phosphate esters, acyl sarcosinates and fatty acid/protein condensates, and mixtures thereof.
21. A composition according to any of Claims 1 to 20 wherein the anionic surfactant comprises an ethoxylated C_8 - C_{22} alkyl sulfate.


WO 96/37595

PCT/US96/07510


32

22. A composition according to any of Claims 1 to 21 wherein the amphoteric surfactant is selected from:

- (a) imidazolinium derivatives of formula (II)



wherein R_1 is C7-C22 alkyl or alkenyl, R_2 is hydrogen or CH_2Z , each Z is independently CO_2M or CH_2CO_2M , and M is H, alkali metal, alkaline earth metal, ammonium or alkanolammonium; and/or ammonium derivatives of formula (III)




wherein R_1 , R_2 and Z are as defined above:

- (b) aminoalkanoates of formula (IV)



and iminodialkanoates of formula (V)



wherein n and m are numbers from 1 to 4, and R_1 and M are independently selected from the groups specified in (a) above; and

- (c) mixtures thereof.

WO 96/37595

PCT/US96/07510

33

23. A composition according to Claim 22 wherein the amphoteric surfactant is selected from the imidazolinium derivatives of formula II and/or ammonium derivatives of formula III.
24. A composition according to any of Claims 1 to 23 wherein the zwitterionic surfactant is selected from alkyl betaine, amido betaine, alkyl sultaine and mixtures thereof.
25. A composition according to Claim 3 wherein the weight ratio of anionic surfactant:other nonionic, zwitterionic and/or amphoteric surfactant is in the range from about 8:1 to about 1:2.
26. A composition according to any of Claims 1 to 25 additionally comprising from 0.01% to 5%, preferably from about 0.04% to about 2% and more preferably from 0.05% to 1% by weight of a cationic or nonionic polymeric skin or hair conditioning agent, selected from cationic and nonionic polysaccharides; cationic and nonionic homopolymers and copolymers derived from acrylic and/or methacrylic acid, cationic and nonionic cellulose resins; cationic copolymers of dimethyldiallylammonium chloride and acrylic acid; cationic homopolymers of dimethyldiallylammonium chloride; cationic polyalkylene and ethoxypolyalkylene imines; quaternized silicones, and mixtures thereof.
27. A composition according to any of Claims 1 to 26 additionally comprising moisturiser selected from glycerin, polyethylene glycol, propylene glycol, sorbitol, polyethylene glycol and polypropylene glycol ethers of methyl glucose, sodium pyrrolidone carboxylic acid, lactic acid, L-proline and mixtures thereof.
28. A composition according to any of Claims 1 to 27 additionally comprising an adduct prepared from vegetable oils containing non-
conjugated polyunsaturated fatty acid esters which are conjugated and elaidinized and then modified via Diels-Alder addition with a member of the group consisting of acrylic acid, fumaric acid and maleic anhydride.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/07510

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) : C11D 3/22, 3/37, 17/00. US CL : 510/119, 135-138, 159, 417, 421-424, 470, 476. According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 510/119, 130, 135-138, 417, 422-424, 470, 476. Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) APS, CAS ONLINE.		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US, A, 4,368,213 (HOLLENBACH ET AL.) 11 January 1983, see col. 2, l. 61 - col. 3, l. 20; col. 4, l. 58 - col. 5, l. 14; col. 5, l. 57 - col. 6, l. 3; the table at the top of col. 7; Examples I-IV.	1-3, 13, 17-19 & 25
X	US, A, 4,446,165 (ROBERTS) 01 May 1984, see col. 3, ll. 3-24, 44-50; col. 4, l. 63 - col. 5, l. 13; col. 8, l. 67 - col. 9, l. 68; col. 10, ll. 1-6; col. 11, l. 24 - col. 12, l. 2.	1-3, 13, 17-19 & 25
X	US, A, 5,160,738 (MACAULEY ET AL.) 03 November 1992, see abstract; col. 3, l. 64 - col. 4, l. 14; col. 4, l. 60 - col. 5, l. 10, especially col. 5, ll. 1 & 2.	1-3, 13, 17-19 & 25
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *Z* document member of the same patent family		
Date of the actual completion of the international search 09 JULY 1996		Date of mailing of the international search report 27 SEP 1996
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer <i>N. Pandey Roy</i> for ARDITH HERTZOG Telephone No. (703) 308-0661

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/07510

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 4-12, 14-16, 20-24 & 26-28
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.